

On the existence and coercive estimates of solutions to the Dirichlet problem for a class of third-order differential equations

A.O. Suleimbekova*, B.M. Musilimov

*M.Kh. Dulaty Taraz Regional University, Taraz, Kazakhstan
(E-mail: suleimbekovaa@mail.ru, bilanmus45@mail.ru)*

As you know, the third order partial differential equation is one of the basic equations of wave theory. For example, in particular, a linearized Korteweg-de Vries type equation with variable coefficients models ion-acoustic waves into plasma and acoustic waves on a crystal lattice. In this paper, the properties of solutions of a class of the third order degenerate partial differential equations with variable coefficients given in a rectangle were studied. Sufficient conditions for the existence and uniqueness of a strong solution have been established. Note that the solution of the degenerate equation does not retain its smoothness, therefore, these difficulties in turn affect the coercive estimates.

Keywords: resolvent, third order differential equation, Dirichlet problem, coercive estimates.

2020 Mathematics Subject Classification: 39A14.

Introduction

In the rectangle $\bar{\Omega} = \{(x, y) : -\pi \leq x \leq \pi; 0 \leq y \leq 1\}$, the problem

$$Lu + \mu u = -k(y) \frac{\partial^3 u}{\partial x^3} - \frac{\partial u^2}{\partial y^2} + a(y) \frac{\partial u}{\partial x} + c(y)u + \mu u = f(x, y) \in L_2(\Omega), \quad (1)$$

$$u_x^{(\alpha)}(-\pi, y) = u_x^{(\alpha)}(\pi, y), \quad \alpha = 0, 1, 2, \quad (2)$$

$$u(x, 0) = u(x, 1), \quad (3)$$

is considered.

Suppose that the coefficients $k(y)$, $a(y)$, $c(y)$ of equation (1) satisfy the conditions:

- 1) $k(y) \geq 0$ is a piecewise continuous function on the segment $[0, 1]$ and $k(0) = 0$;
- 2) $a(y) \geq \delta_0$, $c(y) \geq \delta > 0$ are continuous functions on the segment $[0, 1]$.

Equation (1) degenerates along the line $y = 0$, i.e. at these points equation (1) changes order. This means that solutions do not retain their smoothness, hence these difficulties in turn affect the coercive estimates of solutions.

Many papers [1–13] and the works cited there are devoted to the study of partial differential equations of the third order. From these works and from a review of literary sources, it follows that previously differential equations without degeneracy were mainly studied.

To present the results obtained regarding this work, we will need the following designations and definitions. By $W_2^1(\Omega)$ we denote the S.L. Sobolev space with norm $\|u\|_{2,1,\Omega} = [\|u_y\|_2^2 + \|u_x\|_2^2 + \|u\|_2^2]^{\frac{1}{2}}$. $C_{0,\pi}^\infty(\bar{\Omega})$ is a set consisting of infinitely differentiable functions and satisfying conditions (2)–(3).

*Corresponding author. E-mail: suleimbekovaa@mail.ru

This paper was supported by the grant AP19676466 of the Ministry of Science and Higher Education of Republic of Kazakhstan.

Received: 25 July 2023; Accepted: 29 February 2024.

Definition 1. A function $u(x, y) \in L_2(\Omega)$ is called a strong solution to problem (1)–(3) if there exists a sequence of functions $\{u_n\} \subset C_{0,\pi}^\infty(\overline{\Omega})$, such that

$$\|u_n - u\|_{L_2(\Omega)} \rightarrow 0, \quad \|Lu_n - f\|_{L_2(\Omega)} \rightarrow 0 \text{ as } n \rightarrow 0.$$

Theorem 1. Let the conditions 1) - 2) be fulfilled. Then for $\mu \geq 0$, for any $f(x, y) \in L_2(\Omega)$ there is a unique strong solution to the problem (1)–(3).

Theorem 2. Let conditions 1) - 2) be fulfilled. Then for $\mu \geq 0$, for any $f(x, y) \in L_2(\Omega)$ there is a unique strong solution to the problem(1)–(3) such that the coercive estimate

$$\|u\|_{1,2,\Omega} \leq C \|(L + \mu I)u\|_2$$

is valid for it, where $C > 0$ is a constant, $\|\cdot\|_2$ is the norm of $L_2(\Omega)$.

1 Proof of Theorems 1-2

In what follows, we denote by $(L + \mu I)$ the operator corresponding to problem (1)–(3).

Lemma 1. Let the conditions 1) - 2) be fulfilled. Then the following inequality

$$\|(L + \mu I)u\|_2 \geq (\delta_0 + \lambda) \|u\|_2, \quad (4)$$

holds for all $u \in D(L)$, where $\delta_0 > 0$, $\mu \geq 0$. $D(L)$ is the domain of definition of the operator L .

Proof. Consider the functionality $\langle (L + \mu I)u, u \rangle$, $u \in D(L)$, where $\langle \cdot, \cdot \rangle$ is scalar product in $L_2(\Omega)$. Integrating in parts, we get an estimate (4). Lemma 1 is proved.

Using the Fourier method, we reduce the problem (1)–(3) to the study of the following differential operator with the parameter n ($n = \pm 0, \pm 1, \pm 2, \dots$):

$$(l_n + \mu I)z(y) = -z'(y) + (-ik(y)n^3 + ina(y) + c(y) + \mu)z(y),$$

where $z(y) \in D(l_n)$, $D(l_n)$ is the domain of definition of the operator l_n .

Lemma 2. Let the conditions 1) - 2) be fulfilled. Then the following inequality

$$\|(l_n + \mu I)z\|_2 \geq (\delta_0 + \mu) \|z\|_2$$

holds for all $z(y) \in D(l_n + \mu I)$, where $\|\cdot\|_2$ is the norm of the Hilbert space $L_2(0, 1)$. $D(l_n)$ is the domain of definition of the operator l_n .

Proof. Let us denote by $C_0^2[0, 1]$ the set consisting of doubly differentiable functions and satisfying condition (3). Let $z(y) \in C_0^2[0, 1]$ and consider the functional

$$\langle (l_n + \mu I)z, z \rangle = \int_0^1 [|z'|^2 + (c(y) + \mu)|z|^2 + (in^3k(y) + ina(y))|z|^2] dy. \quad (5)$$

Hence, using the properties of complex numbers, we find that

$$|\langle (l_n + \mu I)z, z \rangle| \geq \int_0^1 [|z'|^2 + (c(y) + \mu)|z|^2] dy \geq \int_0^1 (|z'|^2 + (\delta + \mu)|z|^2) dy. \quad (6)$$

From the last inequality, using the Cauchy-Bunyakovsky inequality, we have

$$\|(l_n + \mu I)z\|_2 \geq (\delta_0 + \mu) \|z\|_2.$$

Hence, and by virtue of the continuity of the norm in $L_2(0, 1)$, we will be convinced of the validity of the last estimate for all $z(y) \in D(l_n)$. Lemma 2 is proved.

Lemma 3. Let the conditions 1) - 2) be fulfilled and $\mu \geq 0$. Then for the operator $(l_n + \mu I)$ there is a bounded inverse operator $(l_n + \mu I)^{-1}$ defined on the whole $L_2(0, 1)$.

Proof. Lemma 3 is also proved as Lemma 2.3 of [14, 15].

Lemma 4. Let the conditions 1) - 2) be fulfilled and $\mu \geq 0$. Then the following estimates are valid for operators $(l_n + \mu I)^{-1}$ and $\frac{d}{dy}(l_n + \mu I)^{-1}$:

$$\|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2} \leq \frac{1}{\delta + \mu}; \quad (7)$$

$$\|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2} \leq \frac{1}{|n| \cdot \delta_0}, \quad n \neq 0; \quad (8)$$

$$\left\| \frac{d}{dy}(l_n + \mu I)^{-1} \right\|_{2 \rightarrow 2} \leq \frac{1}{(\delta + \mu)^{\frac{1}{2}}}, \quad (9)$$

where $\|\cdot\|_{2 \rightarrow 2}$ is the norm of the operator from $L_2(\Omega)$ to $L_2(\Omega)$.

Proof. From Lemma 2 we have

$$\|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2} \leq \frac{1}{\delta + \mu}.$$

Inequality (7) is proved.

Using inequality (5) and properties of complex numbers, we find that

$$\langle (l_n + \mu I)z, z \rangle \geq \left| \int_0^1 (in^3 k(y) + ina(y)) |z|^2 dy \right|. \quad (10)$$

Note that by virtue of condition 1) - 2) the functions $k(y)$ and $a(y)$ do not change signs, therefore, from the inequality (10) we find that

$$|\langle (l_n + \mu I)z, z \rangle| \geq \int_0^1 |in^3 k(y) + ina(y)| \cdot |z|^2 dy. \quad (11)$$

From (11) and given $a(y) \geq \delta_0 > 0$ we have

$$\|(l_n + \mu I)z\|_2 \geq |n| \delta_0 \|z\|_2.$$

Hence, using the definition of the operator norm, we obtain the following estimate:

$$\|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2} \leq \frac{1}{|n| \cdot \delta_0}, \quad n \neq 0.$$

Inequality (8) is proved.

Using inequalities (4) and (6) we find that

$$\frac{1}{\delta + \mu} \|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2} \geq \|z'\|_2^2.$$

Hence, according to the definition of the operator norm, we find

$$\left\| \frac{d}{dy}(l_n + \mu I)^{-1} \right\|_{2 \rightarrow 2} \leq \frac{1}{(\delta + \mu)^{\frac{1}{2}}}.$$

Inequality (9) is proved. Lemma 4 is proved.

Proof of Theorem 1. Using Lemma 3, we obtain that

$$u_k(x, y) = \sum_{n=-k}^{n=k} (l_n + \mu I)^{-1} f_n(y) \cdot e^{inx}$$

is the solution of the following problem:

$$(L + \mu I) u_k(x, y) = f_k(x, y), \quad (12)$$

$$u_{k,x}^{(\alpha)}(-\pi, y) = u_{k,x}^{(\alpha)}(\pi, y), \quad \alpha = 0, 1, 2, \quad (13)$$

$$u_k(x, 0) = u_k(x, 1) = 0, \quad (14)$$

where $f_k(x, y) \rightarrow f(x, y)$, $f_k(x, y) = \sum_{n=-k}^k f_n(y) \cdot e^{inx}$, $i^2 = -1$.

From inequality (4) and using the fundamentality of the sequence $\{f_k(x, y)\}$, we have

$$\|u_k(x, y) + u_m(x, y)\|_2 \leq \frac{1}{\delta + \mu} \|f_k(x, y) - f_m(x, y)\|_2 \rightarrow 0, \quad \text{as } k, m \rightarrow \infty.$$

From the last inequality and by virtue of the completeness of the Hilbert space $L_2(\Omega)$ we have

$$u_k(x, y) \xrightarrow{L_2(\Omega)} u(x, y). \quad (15)$$

Further, using the equalities (12)–(15) for any $f(x, y) \in L_2(\Omega)$, we obtain that

$$u(x, y) = (L + \mu I)^{-1} f = \sum_{n=-\infty}^{n=\infty} (l_n + \mu I)^{-1} f_n(y) \cdot e^{inx} \quad (16)$$

is a strong solution to the problem (1)–(3). The theorem is proved.

Proof of Theorem 2. From (16) by virtue of the orthonormality of the system $\{e^{inx}\}$

$$\begin{aligned} \|u\|_2^2 &= \left\| \sum_{n=-\infty}^{\infty} (l_n + \mu I)^{-1} f_n(y) \cdot e^{inx} \right\|_{L_2(0,1)}^2 = 2\pi \sum_{n=-\infty}^{\infty} \|(l_n + \mu I)^{-1} f_n(y) \cdot e^{inx}\|_{L_2(0,1)}^2 \leq \\ &\leq 2\pi \sum_{n=-\infty}^{\infty} \|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 \cdot \|f_n(y)\|_{L_2(0,1)}^2 \leq \sup_{\{n\}} \|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 2\pi \cdot \sum_{n=-\infty}^{\infty} \|f_n(y)\|_{L_2(0,1)}^2 \leq \\ &\leq \sup_{\{n\}} \|(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 \cdot \|f(x, y)\|_{L_2(\Omega)}^2. \end{aligned} \quad (17)$$

Here we note that we used by virtue of the orthonormality of the system $\{e^{inx}\}$, i.e.

$$\|f(x, y)\|_2^2 = \left\| \sum_{n=-\infty}^{\infty} f_n(y) \cdot e^{inx} \right\|_{L_2(\Omega)}^2 = 2\pi \cdot \sum_{n=-\infty}^{\infty} \|f_n(y)\|_{L_2(0,1)}^2.$$

From estimates (17), (4) and (7) we obtain that

$$\|u\|_{L_2(\Omega)}^2 \leq \left(\frac{1}{\delta + \mu}\right)^2 \|f(x, y)\|_{L_2(\Omega)}^2.$$

From here we finally have

$$\|u\|_2 \leq \|f(x, y)\|_{L_2(\Omega)}, \quad (18)$$

where $C_1 = \frac{1}{\delta+\mu}$.

Next, we calculate the norm $\|u_x\|_2$:

$$\begin{aligned} \|u_x\|_{L_2(\Omega)}^2 &= \left\| \sum_{n=-\infty}^{\infty} in(l_n + \mu I)^{-1} e^{inx} \right\|_{L_2(\Omega)}^2 \leq \sup_{\{n\}} \|in(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 \cdot 2\pi \cdot \sum_{n=-\infty}^{\infty} \|f_n(y)\|_{L_2(0,1)}^2 \leq \\ &\leq \sup_{\{n\}} |n|^2 \cdot \|in(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 \cdot \|f(x, y)\|_{L_2(\Omega)}^2. \end{aligned}$$

Hence and from inequality (8) we have

$$\|u_x\|_{L_2(\Omega)}^2 \leq \sup_{\{n\}} |n|^2 \|in(l_n + \mu I)^{-1}\|_{2 \rightarrow 2}^2 \cdot \|f(x, y)\|_{L_2(\Omega)}^2 \leq \sup_{\{n\}} |n|^2 \cdot \frac{1}{|n|^2 \cdot \delta_0^2} \|f(x, y)\|_{L_2(\Omega)}^2.$$

Hence

$$\|u_x\|_{L_2(\Omega)} \leq C_2 \|f(x, y)\|_{L_2(\Omega)}, \quad (19)$$

where $C_2 = \frac{1}{(\delta+\mu)^{\frac{1}{2}}}$.

Then, repeating the above calculations, we get the following estimate

$$\|u_y\|_{L_2(\Omega)} \leq C_3 \|f(x, y)\|_{L_2(\Omega)}, \quad (20)$$

where $C_2 = \frac{1}{\delta_0}$.

Using the equalities (18)–(20), we find that

$$\|u\|_{2,1,\Omega} \leq C \|f(x, y)\|_{L_2(\Omega)},$$

where $C = \max\{C_1, C_2, C_3\}$. The theorem is proved.

Acknowledgments

This paper was supported by the grant AP19676466 of the Ministry of Science and Higher Education of Republic of Kazakhstan.

Author Contributions

A.O. Suleimbekova collected and analyzed data, and led manuscript preparation. B.M. Musilimov assisted in data collection and analysis. All authors participated in the revision of the manuscript and approved the final submission.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Ashyralyev A. On the stability of nonlocal boundary value problem for a third order PDE / A. Ashyralyev, K. Belakroum // Third International Conference Of Mathematical Sciences. — 2019. — Vol. 2183, No. 1. — P. 1–4. <https://doi.org/10.1063/1.5136174>
- 2 Ashyralyev A. On stability of the third order partial delay differential equation with involution and Dirichlet condition / A. Ashyralyev, S. Ibrahim, E. Hincal // Bulletin of The Karaganda University. Mathematics series. — 2021. — No. 2(102). — P. 25–34. <https://doi.org/10.31489/2021M2/25-34>
- 3 Maher A. On the cases of explicit solvability of a third order partial differential equation / A. Maher, Ye.A. Utkina // Iranian journal of science and technology transaction a science. — 2010. — Vol. 34, No. 2. — P. 103–112. <https://doi.org/10.22099/IJSTS.2010.2169>
- 4 Saut J.C. Remarks on the Korteweg-de Vries equation / J.C. Saut, R. Temam // Israel Journal of Mathematics. — 1976. — Vol. 24, No. 1. — P. 78–87. <https://doi.org/10.1007/BF02761431>
- 5 Belakroum K. A note on nonlocal boundary value problem for a third-order partial differential equation / K. Belakroum, A. Ashyralyev, A. Guezane-Lakoud // Filomat. — 2018. — Vol. 32, No. 3. — P. 801–808. <https://doi.org/10.2298/FIL1803801B>
- 6 Gao P. Stochastic Korteweg-de Vries equation in a bounded domain / P. Gao // App Math and Computation. — 2017. — Vol. 310. — P. 97–111. <https://doi.org/10.1016/j.amc.2017.04.031>
- 7 Xiang S. Zero controllability of the linearized Korteweg-De Vries equation using the reverse control approach / S. Xiang // Siam Journal on control and optimization. — 2019. — Vol. 57, No. 2. — P. 1493–1515. <https://doi.org/10.1137/17M1115253>
- 8 Aliyev Z.S. On some nonlocal inverse boundary problem for partial differential equations of third order / Z.S. Aliyev, Y.T. Mehraliyev, E.H. Yusifova // Turkish Journal of mathematics. — 2021. — Vol. 45, No. 4. — P. 1871–1886. <https://doi.org/10.3906/mat-2101-37>
- 9 Лукина Г.А. Краевые задачи с интегральными граничными условиями для линеаризованного уравнения Кортевега-де Фриза / Г.А. Лукина // Вестн. ЮУрГУ. Сер. Мат. моделир. и программир. — 2011. — Т. 234, № 17. — С. 52–61.
- 10 Dzhuraev T.D. On the Goursat and Dirichlet problems for equation of the third-order / T.D. Dzhuraev, O.S. Zikirov // Nonlinear Oscillations. — 2008. — Vol. 11, No. 3. — P. 320–330. <https://doi.org/10.1007/s11072-009-0033-0>
- 11 Muratbekov M.B. Coercive solvability of odd-order differential equation and its applications / M.B. Muratbekov, M.M. Muratbekov, K.N. Ospanov // Doklady Mathematics. — 2010. — Vol. 82, No. 3. — P. 1–3. <https://doi.org/10.1134/S1064562410060189>
- 12 Muratbekov M.B. Separability and estimates of the cross-sections of sets associated with the domain of definition of a nonlinear operator of the Schrodinger type / M.B. Muratbekov // Differential equations. — 1991. — Vol. 27, No. 6. — P. 1034–1042.
- 13 Muratbekov M.B. One-dimensional Schrodinger operator with a negative parameter and its applications to the study of the approximation numbers of a singular hyperbolic operator / M.B. Muratbekov, M.M. Muratbekov // Filomat. — 2018. — Vol. 32, No. 3. — P. 785–790. <https://doi.org/10.2298/FIL1803785M>
- 14 Muratbekov M.B. Existence, compactness, estimates of eigenvalues and s-numbers of a resolvent for a linear singular operator of the korteweg-de vries type / M.B. Muratbekov, A.O. Suleimbekova // Filomat. — 2022. — Vol. 36, No. 11. — P. 3691–3702. <https://doi.org/10.2298/FIL2211691M>
- 15 Suleimbekova A.O. Separability of the third-order differential operator given on the whole plane / A.O. Suleimbekova // Bulletin of the Karaganda University. Mathematics series. — 2022. — Vol. 105, No. 1. — P. 109–117. <https://doi.org/10.31489/2022M1/109-117>

Үшінші ретті дифференциалдық теңдеулердің бір класы үшін Дирихле есебі шешімдерінің бар болуы және коэрцитивті бағалаулары туралы

А.О. Сулеймекова, Б.М. Мусилимов

M.X. Dulatov atyndaǵы Taraz ənqırlık universiteti, Taraz, Қазақстан

Білетініміздей үшінші ретті дербес туындылы дифференциалдық теңдеулер толқындар теориясының негізгі теңдеулерінің бірі. Мысалы, айнымалы коэффициентті сызықталған Кортевег—де Фриз типті теңдеуі иондық акустикалық толқындарды кристалдық тордағы плазмалық және акустикалық толқындарға модельдейді. Жұмыста тіктөртбұрышта берілген айнымалы коэффициентті үшінші ретті дербес туындылы еселенген теңдеулердің бір класының шешімдерінің қасиеттері зерттелген. Күшті шешімнің бар болуы мен жалғыздығына жеткілікті шарттар алынған. Еселенген теңдеудің шешімі өзінің тегістігін сактамайтынын ескерсек, бұл қындықтар өз кезегінде коэрцитивті бағалауга әсер етеді.

Кілт сөздер: резольвента, үшінші ретті дифференциалдық теңдеулер, Дирихле есебі, коэрцитивті бағалаулар.

О существовании и коэрцитивных оценках решений задачи Дирихле для одного класса дифференциальных уравнений третьего порядка

А.О. Сулеймекова, Б.М. Мусилимов

Taraazskiy regionalnyy universitet imeni M.X. Dulatov, Taraaz, Kazahstan

Как известно, уравнения в частных производных третьего порядка являются одним из основных уравнений теории волн. В частности, линеаризованное уравнение типа Кортевега—де Фриза с переменными коэффициентами моделирует ионно-акустические волны в плазменные и акустические волны на кристаллической решетке. В данной работе исследованы свойства решений одного класса вырождающихся уравнений в частных производных третьего порядка с переменными коэффициентами, заданных в прямоугольнике. Установлены достаточные условия существования и единственности сильного решения. Заметим, что решение вырождающегося уравнения не сохраняет свою гладкость, следовательно, эти трудности, в свою очередь, влияют на коэрцитивные оценки.

Ключевые слова: резольвента, дифференциальные уравнения третьего порядка, задача Дирихле, коэрцитивные оценки.

References

- 1 Ashyralyev, A., & Belakroum, K. (2019). On the stability of nonlocal boundary value problem for a third order PDE. *Third International Conference Of Mathematical Sciences*, 2183(1), 1–4. <https://doi.org/10.1063/1.5136174>
- 2 Ashyralyev, A., Ibrahim, S., & Hincal, E. (2021). On stability of the third order partial delay differential equation with involution and Dirichlet condition. *Bulletin of the Karaganda University. Mathematics series*, 2(102), 25–34. <https://doi.org/10.31489/2021M2/25-34>
- 3 Maher, A., & Utkina, Ye.A. (2010). On the cases of explicit solvability of a third order partial differential equation. *Iranian journal of science and technology transaction a science*, 34(2), 103–112. <https://doi.org/10.22099/IJSTS.2010.2169>

- 4 Saut, J.C., & Temam, R. (1976). Remarks on the Korteweg-de Vries equation. *Israel Journal of Mathematics*, 24(1), 78–87. <https://doi.org/10.1007/BF02761431>
- 5 Belakroum, K., Ashyralyev, A., & Guezane-Lakoud A. (2018). A note on nonlocal boundary value problem for a third-order partial differential equation. *Filomat*, 32 (3), 801–808. <https://doi.org/10.2298/FIL1803801B>
- 6 Gao, P. (2017). Stochastic Korteweg-de Vries equation in a bounded domain. *App Math and Computation*, 310, 97–111. <https://doi.org/10.1016/j.amc.2017.04.031>
- 7 Xiang, S.(2019). Zero controllability of the linearized Korteweg-De Vries equation using the reverse control approach. *Siam Journal on control and optimization*, 57(2), 1493–1515. <https://doi.org/10.1137/17M1115253>
- 8 Aliyev, Z.S., Mehraliyev, Y.T., & Yusifova, E.H. (2021). On some nonlocal inverse boundary problem for partial differential equations of third order. *Turkish Journal of mathematics*, 45(4), 1871–1886. <https://doi.org/10.3906/mat-2101-37>
- 9 Lukina, G.A. (2011). Kraevye zadachi s integralnymi granichnymi usloviami dlia linearizovannogo uravneniiia Kortevega-de Friza [Boundary value problems with integral boundary conditions for the linearized Korteweg-de Vries equation]. *Vestnik Yuzhno-Uralskogo gosudarstvennogo universiteta. Seria Matematicheskoe modelirovanie i programmirovaniye — Bulletin of SUSU. Ser. Math. modeling and programming*, 234 17, 52–61 [in Russian].
- 10 Dzhuraev, T.D., & Zikirov, O.S. (2008). On the Goursat and Dirichlet problems for equation of the third-order. *Nonlinear Oscillations*, 11(3), 320–330. <https://doi.org/10.1007/s11072-009-0033-0>
- 11 Muratbekov, M.B., Muratbekov, M.M., & Ospanov, K.N (2010). Coercive solvability of odd-order differential equation and its applications. *Doklady Mathematics*, 82(3), 1–3. <https://doi.org/10.1134/S1064562410060189>
- 12 Muratbekov, M.B. (1991). Separability and estimates of the cross-sections of sets associated with the domain of definition of a nonlinear operator of the Schrodinger type. *Differential equation*, 27(6), 1034–1042.
- 13 Muratbekov, M.B., & Muratbekov, M.M. (2018). One-dimensional Schrodinger operator with a negative parameter and its applications to the study of the approximation numbers of a singular hyperbolic operator. *Filomat*, 32(3), 785–790. <https://doi.org/10.2298/FIL1803785M>
- 14 Muratbekov, M.B., & Suleimbekova, A.O. (2022). Existence, compactness, estimates of eigenvalues and s-numbers of a resolvent for a linear singular operator of the korteweg-de vries type. *Filomat*, 36(11), 3691–3702. <https://doi.org/10.2298/FIL2211691M>
- 15 Suleimbekova, A.O. (2022). Separability of the third-order differential operator given on the whole plane. *Bulletin of the Karaganda University. Mathematics series*, 1(105), 109–117. <https://doi.org/10.31489/2022M1/109-117>

*Author Information**

Ainash Ospanovna Suleimbekova (*corresponding author*) — Doctor of philosophy (PhD), Department of Mathematics, M.Kh. Dulaty Taraz Regional University, 7 Suleymenov street, Taraz, 080000, Kazakhstan; e-mail: suleimbekova@mail.ru; <https://orcid.org/0000-0003-1865-4822>

Bilibai Musilimovich Musilimov — Candidate of physical and mathematical sciences, Associate professor, Department of Mathematics, M.Kh. Dulaty Taraz Regional University, 7 Suleymenov street, Taraz, 080000, Kazakhstan; e-mail: bilanmus45@mail.ru

*The author's name is presented in the order: First, Middle and Last Names.