

S. H. Salim<sup>1</sup>, K. H.F. Jwamer<sup>2,\*</sup>, R. K. Saeed<sup>3</sup><sup>1</sup>College of Basic Education, University of Sulaimani, Sulaymaniyah, Iraq;<sup>2</sup>College of Science, University of Sulaimani, Sulaymaniyah, Iraq;<sup>3</sup>College of Science, Salahaddin University, Erbil, Iraq

(E-mail: sarfraz.salim@univsul.edu.iq, karwan.jwamer@univsul.edu.iq, rostam.saeed@su.edu.krd)

## Solving Volterra-Fredholm integral equations by natural cubic spline function

Using the natural cubic spline function, this paper finds the numerical solution of Volterra-Fredholm integral equations of the second kind. The proposed method is based on employing the natural cubic spline function of the unknown function at an arbitrary point and using the integration method to turn the Volterra-Fredholm integral equation into a system of linear equations concerning to the unknown function. An approximate solution can be easily established by solving the given system. This is accomplished with the help of a computer program that runs on Python 3.9.

*Keywords:* Volterra integral equation, Fredholm integral equation, spline function.

### Introduction

Integral equations can be used to express a variety of mathematical physics topics. Some of these will be examined and treated explicitly as examples. It would be nearly impossible to compile a list of such applications. To say that integral equations play a role in practically every area of applied mathematics and mathematical physics is an understatement; because, the literature on integral equations and their applications is extensive.

Many researches have been conducted in recent years, with the results revealing the interaction of Fredholm integral equation, Volterra integral equation, Volterra-Fredholm integral equation, and numerical solutions of these three types of the integral equation.

The linear Volterra-Fredholm integral equations (VFIEs) of the following type were taken into consideration in this work:

$$u(x) = f(x) + \lambda_1 \int_a^x K(x, t)u(t)dt + \lambda_2 \int_a^b L(x, t)u(t)dt, \quad (1)$$

where the functions  $f(x)$ , and the kernels  $K(x, t)$  and  $L(x, t)$  are known  $L^2$  analytic functions and  $\lambda_1$ ,  $\lambda_2$  are arbitrary constants,  $x$  is variable and  $u(x)$  is the unknown continuous function to be determined. Numerous applications in the fields of physics, fluid dynamics, electrodynamics, and biology include the use of these equations. These integral equations are a reduction of several boundary value formulations using Neumann, Dirichlet, or boundary conditions. Additionally, they offer mathematical models for the spread of an epidemic as well as a host of other physical and biological problems.

Since the analytical solution to VFIEs generally does not exist outside of special cases, the numerical method is the most successful and efficient way to solve these issues. In order to solve VFIEs, a number of numerical and approximative techniques have been developed, including the linear and quadratic spline functions by Salim, et. al. [1, 2], Taylor polynomial by Yalcinbas and Sezer [3]; Yalcinbas [4], the least square method and Chebyshev polynomials by Dastjerdi and Ghaini [5].

---

\*Corresponding author.

E-mail: karwan.jwamer@univsul.edu.iq

Also, Lagrange collocation method by Wang [6], Series solution, successive approximation method and method of successive substitutions by Saeed and Berdawood [7], trigonometric Functions and Laguerre Polynomials by Hasan and Sulaiman [8], Touchard Polynomials (T-Ps) method by Al-Miah and Taie [9]. Some iterative numerical methods by Micula [10], Taylor polynomial by Didgara and Vahidi [11]. For additional information, the reader might turn to the following references and the references given there: (Jerry [12], Atkinson [13], Lange and Herbert [14], Kaminaka and Wadati [15], Ladopoulos [16], Corduneanu [17], Saeed and Aziz [18] and Jaber and Alrammahi [19]).

Equation (1) will be studied in this work using the natural cubic spline function. The rest of this paper is structured as follows. Our approach is introduced in Section 2 for solving Equation (1). We examine various numerical examples proving the viability of our method in Section 3. Some conclusions will be made in Section 4.

### 1 Description of the method

In this section, we solve Equation (1) by using the quadratic spline function (Cheney and Kincaid [20], Saeed *et. al.* [21]). The unknown function  $u(x)$  in (1) is approximated by the quadratic spline function  $S(x)$ . In the interval  $[x_i, x_{i+1}]$ , the quadratic spline function is defined by the following formula:

$$S_i(x) = A_i(x)S_i + B_i(x)S_{i+1} + C_i(x)S'_i + D_i(x)S'_{i+1}, \quad (2)$$

where  $A_i(x) = 1 - 3\frac{(x-t_i)^2}{h^2} + 2\frac{(x-t_i)^3}{h^3}$ ,  $B_i(x) = 1 - A_i(x)$ ,  $C_i(x) = \frac{(x-t_i)(x-t_{i+1})^2}{h^2}$ ,  $D_i(x) = \frac{(x-t_{i+1})(x-t_i)^2}{h^2}$ , and  $h = x_{i+1} - x_i$  for all  $i = 0, 1, \dots, n-1$ . Now substituting (2) in (1) and letting  $x = x_i$ , we get

$$\begin{aligned} S_i &= f_i + \lambda_1 \int_a^{x_i} K(x_i, t)S(t) + \lambda_2 \int_a^b L(x_i, t)S(t)dt \\ &= f(x_i) + \lambda_1 \left[ \sum_{j=0}^{j=i-2} \int_{x_j}^{x_{j+1}} K(x_i, t)[A_j(t)S_j + B_j(t)S_{j+1} + C_j(t)S'_j + D_j(t)S'_{j+1}]dt \right. \\ &\quad \left. + \int_{x_{i-1}}^{x_i} K(x_i, t)[A_{i-1}(t)S_{i-1} + B_{i-1}(t)S_i + C_i(t)S'_i + D_i(t)S'_i]dt \right] \\ &\quad + \lambda_2 \left[ \int_{x_0=a}^{x_1} L(x_i, t)S_0(t)dt + \int_{x_1}^{x_2} L(x_i, t)S_1(t)dt + \dots + \int_{x_{n-1}}^{x_n=b} L(x_i, t)S_{n-1}(t)dt \right] \\ &= f(x_i) + \lambda_1 \left[ \sum_{j=0}^{j=i-2} \int_{x_j}^{x_{j+1}} K(x_i, t)[A_j(t)S_j + B_j(t)S_{j+1} + C_j(t)S'_j + D_j(t)S'_{j+1}]dt \right. \\ &\quad \left. + \int_{x_{i-1}}^{x_i} K(x_i, t)[A_{i-1}(t)S_{i-1} + B_{i-1}(t)S_i + C_i(t)S'_i + D_i(t)S'_i]dt \right] \\ &\quad + \lambda_2 \left[ \int_{x_0=a}^{x_1} L(x_i, t)[A_0(t)S_0 + B_0(t)S_1 + C_0(t)S'_0 + D_0(t)S'_1]dt \right. \\ &\quad + \int_{x_1}^{x_2} L(x_i, t)[A_1(t)S_1 + B_1(t)S_2 + C_1(t)S'_1 + D_1(t)S'_2]dt + \dots \\ &\quad \left. + \int_{x_{n-1}}^{x_n=b} L(x_i, t)[A_{n-1}(t)S_{n-1} + B_{n-1}(t)S_n + C_{n-1}(t)S'_{n-1} + D_{n-1}(t)S'_n]dt \right]. \end{aligned}$$

By computing the integrals in the above equation using the trapezoidal rule, we get

$$S_i = f_i + \frac{h}{2}(\lambda_1 K_{i0} + \lambda_2 L_{i0})S_0 + h \sum_{j=1}^{i-1} (\lambda_1 K_{ij} + \lambda_2 L_{ij})S_j + \frac{h}{2}(\lambda_1 K_{ii} + \lambda_2 L_{ii})S_i, \quad (3)$$

for  $i = 0, 1, \dots, n$ .

In this way, Equation (3) constructs a system of linear equations concerning to the unknown function  $S_i$ . Briefly, this system can be rewritten as follows:

$$CS = F, \quad (4)$$

where

$$S = \begin{bmatrix} S_0 \\ S_1 \\ \vdots \\ S_n \end{bmatrix}, \quad F = \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_n \end{bmatrix}, \quad C = [C_0 \quad C_1 \quad C_2 \quad \dots \quad C_{n-1} \quad C_n],$$

$$C_0 = \begin{bmatrix} 1 - \frac{\lambda_2 h}{2} L_{00} \\ -\frac{h}{2}(2\lambda_1 K_{10} + \lambda_2 L_{10}) \\ -\frac{h}{2}(\lambda_1 K_{20} + \lambda_2 L_{20}) \\ -\frac{h}{2}(\lambda_1 K_{30} + \lambda_2 L_{30}) \\ \vdots \\ -\frac{h}{2}(\lambda_1 K_{n0} + \lambda_2 L_{n0}) \end{bmatrix}, \quad C_1 = \begin{bmatrix} -\lambda_2 h L_{01} \\ 1 - (\frac{h}{2}(\lambda_1 K_{11} + 2\lambda_2 L_{11})) \\ -\frac{h}{2}(3\lambda_1 K_{21} + 2\lambda_2 L_{21}) \\ -h(\lambda_1 K_{31} + \lambda_2 L_{31}) \\ \vdots \\ -h(\lambda_1 K_{n1} + \lambda_2 L_{n1}) \end{bmatrix},$$

$$C_2 = \begin{bmatrix} -\lambda_2 h L_{02} \\ -\frac{h}{2}(2\lambda_2 L_{12} - \lambda_1 K_{10}) \\ 1 - \frac{h}{2}(\lambda_1 K_{22} + 2\lambda_2 L_{22}) \\ -\frac{h}{2}(3\lambda_1 K_{32} + 2\lambda_2 L_{32}) \\ \vdots \\ -h(\lambda_1 K_{n2} + \lambda_2 L_{n2}) \end{bmatrix}, \quad \dots, \quad C_{n-1} = \begin{bmatrix} -\lambda_2 h L_{0(n-1)} \\ -\lambda_2 h L_{1(n-1)} \\ -\lambda_2 h L_{2(n-1)} \\ \vdots \\ 1 - \frac{h}{2}(\lambda_1 K_{(n-1)(n-1)} + 2\lambda_2 L_{(n-1)(n-1)}) \\ -\frac{h}{2}(\lambda_1 K_{n(n-1)} + 2\lambda_2 L_{n(n-1)}) \end{bmatrix},$$

and

$$C_n = \begin{bmatrix} -\frac{\lambda_2 h}{2} L_{0n} \\ -\frac{\lambda_2 h}{2} L_{1n} \\ -\frac{\lambda_2 h}{2} L_{2n} \\ \vdots \\ -\frac{\lambda_2 h}{2} L_{(n-1)(n-1)} \\ 1 - \frac{h}{2}(\lambda_1(K_{nn} + \lambda_2 L_{nn})) \end{bmatrix}.$$

In the sequel, using a standard rule to the resulting system yields an approximate solution of Equation (1) as  $S_i(x)$  given by Equation (2).

## 2 Numerical examples

In this section, we present three examples to illustrate the efficiency and accuracy of the proposed method. The computed errors  $e_i$  are defined by  $e_i = |u_i - S_i|$ , where  $u_i$  is the exact solution of Equation

(1) and  $S_i$  is an approximate solution of the same equation. Also we compute Least square error (LSE) which is defined by formula  $= \sum_{i=0}^n (u_i - S_i)^2$  and all computations are performed using the Python program.

*Example 1.* Consider the linear Volterra-Fredholm integral equation

$$u(x) = -\frac{x^2}{2} - \frac{7x}{2} + 2 + \int_0^x u(t)dt + \int_0^1 xu(t)dt.$$

The exact solution to this equation is given by  $u(x) = x + 2$ .

Table (1) demonstrates LSE obtained from applying our method to Example (1) for  $n = 5$ .

Table 1

The Numerical Results for Example (1) with  $n = 5$

$x_i$	$u_i$	$S_i$	$ u_i - S_i $	$ u_i - S_i ^2$
0	2	2	0	0
0.2	2.2	2.167670	0.323298	0.00104522
0.4	2.4	2.369779	0.302205	0.00091328
0.6	2.6	2.533617	0.06738252	0.00454041
0.8	2.8	3.0875551	0.2875513	0.08268795
1	3	3.0788111	0.0788111	0.00621119
LSE				$9.5398050 \times 10^{-2}$

*Example 2.* Consider the linear Volterra-Fredholm integral equation

$$u(x) = 2\cos(x) - 1 + \int_0^x (x - t)u(t)dt + \int_0^\pi u(t)dt.$$

The exact solution to this equation is given by  $u(x) = \cos(x)$ .

Table (2) demonstrates LSE obtained from applying our method to Example (2) for  $n = 5$ .

Table 2

The Numerical Results for Example (2) with  $n = 5$

$x_i$	$u_i$	$S_i$	$ u_i - S_i $	$ u_i - S_i ^2$
0	1	0.9020653	0.0979346	$9.59119752 \times 10^{-3}$
$\frac{\pi}{5}$	0.8090169	0.80093654	0.00808045	$6.52937193 \times 10^{-5}$
$\frac{2\pi}{5}$	0.3090169	0.38139274	0.07237575	$5.23824911 \times 10^{-3}$
$\frac{3\pi}{5}$	-0.3090169	-0.15642311	0.15259389	$2.32848948 \times 10^{-2}$
$\frac{4\pi}{5}$	-0.8090169	0.8466296	0.0376126	$1.41470781 \times 10^{-3}$
$\pi$	-1.	-0.9488828	0.0511172	$2.61296767 \times 10^{-3}$
LSE				$4.22073106838 \times 10^{-2}$

*Example 3.* Consider the linear Volterra-Fredholm integral equation

$$u(x) = -\frac{x^5}{10} + 2x^3 - \frac{x^2}{2} - \frac{3x}{2} + \frac{1}{10} + \int_0^x (x + t)u(t)dt + \int_0^1 (x - t)u(t)dt.$$

The exact solution to this equation is given by  $u(x) = 2x^3 + 1$ .

Table (3) demonstrates LSE obtained from applying our method to Example (3) for  $n = 5$ .

Table 3

The Numerical Results for Example (3) with  $n = 5$ 

$x_i$	$u_i$	$S_i$	$ u_i - S_i $	$ u_i - S_i ^2$
0	1	0.9320698	0.0679302	0.00461451
0.2	1.016	0.91835798	0.09764202	0.00953396
0.4	1.128	0.99225104	0.13574896	0.01842778
0.6	1.432	1.24719269	0.18480731	0.03415374
0.8	2.024	1.83381163	0.19018837	0.03617162
1	3	2.72882412	0.27117588	0.07353636
LSE				$1.76437975899 \times 10^{-1}$

Table 4

LSE for different values of  $n$  for Examples (1)–(3)

LSEn	10	20	30	40	50
Example 1	$2.02193681 \times 10^{-2}$	$5.21316981 \times 10^{-3}$	$2.36837446 \times 10^{-3}$	$1.348919901 \times 10^{-3}$	$8.7012926 \times 10^{-4}$
Example 2	$3.03911660 \times 10^{-3}$	$1.87835246 \times 10^{-4}$	$3.79028932 \times 10^{-5}$	$1.24158506 \times 10^{-5}$	$5.28945050 \times 10^{-6}$
Example 3	$1.95064939 \times 10^{-2}$	$2.31413450 \times 10^{-3}$	$6.752544397 \times 10^{-4}$	$2.82826976 \times 10^{-4}$	$1.44203795 \times 10^{-4}$

### 3 Conclusion

In this paper the cubic spline function is used to solve linear Volterra-Fredholm integral equations, and it is a powerful numerical approach. The numerical results in the present section demonstrate that the proposed method can successfully tackle the Volterra-Fredholm type problem. Table (4) shows that the proposed method has extremely good stability; as  $n$  increases, the error decreases at first and then stabilizes. We also conclude that we have high accuracy when the exact solution is a trigonometric function. The present method can be easily extended to systems of Volterra-Fredholm integral equations and systems of Volterra-Fredholm integro-differential equations. The current method may be simply extended to Volterra-Fredholm integral equations and Volterra-Fredholm integro-differential equations.

### References

- 1 Salim, H.S., Jwamer, K.H.F., & Saeed, R.K. (2022). Solving Volterra-Fredholm Integral Equation by linear spline function. *Global and Stochastic Analysis*, 9(2), 99–107.
- 2 Salim, H.S., Saeed, R.K., & Jwamer, K.H.F. (2022). Solving Volterra-Fredholm Integral Equation by quadtartic spline function. *Journal of Al-Qadsiyah for computer science and mathematics*, 14(4), 10–19.
- 3 Yalcinbas, S., & Seser, M. (2000). The approximation solution of high-order linear volterra-Fredholm integro-differential equations in terms of Taylor polynomials. *Applied Mathematics and Computation*, 112(2-3), 291–308.
- 4 Yalcinbas, S. (2002). Taylor polynomial solution of nonlinear Volterra-Fredholm integral equations. *Applied Mathematics and Computation*, 127(2-3), 195–206.
- 5 Dastjerdi, H.L., & Ghaini, F.M.M. (2012). Numerical solution of Volterra-Fredholm integral equations by moving least square method and Chebyshev polynomials. *Applied Mathematical Modelling*, 36, 3283–3288.
- 6 Wang, K.Y., & Wang, Q.S. (2013). Lagrange collocation method for solving Volterra-Fredholm integral equations. *Applied Mathematics and Computation*, 219(21), 10434–10440.

- 7 Saeed, R.K., & Berdawood, K.A. (2016). Solving Two-dimensional Linear Volterra-Fredholm Integral Equations of the Second Kind by Using Successive Approximation Method and Method of Successive Substitutions. *ZANCO Journal of Pure and Applied Sciences*, 28(2), 35–46.
- 8 Hasan, P.M.A., & Sulaiman, N.A. (2016). Numerical Treatment of Mixed Volterra-Fredholm Integral Equations Using Trigonometric Functions and Laguerre Polynomials. *ZANCO Journal of Pure and Applied Sciences*, 30(6), 97–106.
- 9 Al-Miah, J.T.A., & Taie, A.H.S. (2019). A new Method for Solutions Volterra-Fredholm Integral Equation of the Second Kind. *IOP Conf. Series: Journal of Physics: Conf. Series*, 1294, 032026.
- 10 Micula, S. (2019). On Some Iterative Numerical Methods for Mixed Volterra-Fredholm Integral Equations. *Symmetry*, 11(10), 1200. <https://doi.org/10.3390/sym11101200>
- 11 Didgara, M., & Vahidi, A. (2020). Approximate Solution of Linear Volterra-Fredholm Integral Equations and Systems of Volterra-Fredholm Integral Equations using Taylor Expansion Method. *Iranian Journal of Mathematical Sciences and Informatics*, 15(1), 31–50.
- 12 Jerry, A.J. (1985). *Introduction to Integral Equation with Application*. Marcel Dekker.
- 13 Atkinson, K.E. (1997). *The numerical solution of integral equation of the second kind*, 4. Cambridge university press.
- 14 Lange, W.A., & Herbert, J.M. (2011). Symmetric versus asymmetric discretization of the integral equations in polarizable continuum solvation models. *Chemical Physics Letters*, 509(1), 77–87.
- 15 Kaminaka, T., & Wadati, M. (2011). Higher order solutions of Lieb-Liniger integral equation. *Physics Letters A*, 375(24), 2460–2464.
- 16 Ladopoulos, E.G. (2011). Reserves exploration by real-time expert seismology and non linear singular integral equations. *Oil Gas and Coal Technol.*, 5(4), 299–315.
- 17 Corduneanu, C. (1991). *Integral Equations and applications*. Cambridge University Press, United Kingdom.
- 18 Saeed, R.K., & Aziz, K.M. (2008). Approximate Solution of the System of Nonlinear Fredholm Integral Equations of the Second Kind Using Spline Function. *Journal of Kirkuk University-Scientific Studies*, 3, 113–128.
- 19 Jaber, K.F., & Alrammahi, A. (2020). Spline Technique for Second Type of Fredholm Integral Equations. *Journal of Kufa for Mathematics and Computer*, 7, 31–40.
- 20 Cheney, W., & Kincaid, D. (1999). *Numerical Mathematics and Computing*. Brooks/Cole Publication Company.
- 21 Saeed, R.K., Jwamer, K.H.F., & Hamasalh, F.K. (2015). *Introduction to Numerical Analysis*. First Edition, University of Sulaimani.

С.Х. Салим<sup>1</sup>, К.Х.Ф. Жвамер<sup>2</sup>, Р.К. Саид<sup>3</sup>

<sup>1</sup> Сулейман университетінің негізгі білім беру колледжі, Сулеймания, Ирак;

<sup>2</sup> Сулейман университетінің жаратылыстану ғылымдары колледжі, Сулеймания, Ирак;

<sup>3</sup> Жаратылыстану ғылымдары колледжі, Салахаддин университеті, Эрбил, Ирак

## Вольтерра-Фредгольм интегралдық теңдеулерін кубтық сплайн-функциясымен шешу

Мақалада табиғи кубтық сплайн-функциясын қолданып екінші текті Вольтерра-Фредгольм аралас интегралдық теңдеулерінің сандық шешімі табылған. Ұсынылған әдіс еркін нүктеде белгісіз функцияның табиғи кубтық сплайн-функциясын қолдануға және Вольтерра-Фредгольм интегралдық теңдеуін

белгісіз функцияға қатысты сызықтық теңдеулер жүйесіне түрлендіру үшін интегралдау әдісін қолдануға негізделген. Бұл жүйені шешу арқылы жуық шешімді табу оңай. Бұған Python 3.9-да жұмыс істейтін компьютерлік бағдарлама арқылы қол жеткізіледі.

*Клт сөздер:* Вольтерра интегралдық теңдеуі, Фредгольм интегралдық теңдеуі, сплайн-функциясы.

С.Х. Салим<sup>1</sup>, К.Х.Ф. Жвамер<sup>2</sup>, Р.К. Саид<sup>3</sup>

<sup>1</sup> Колледж базового образования Сулейманского университета, Сулеймания, Ирак;

<sup>2</sup> Колледж естественных наук Сулейманского университета, Сулеймания, Ирак;

<sup>3</sup> Колледж естественных наук, Университет Салахаддина, Эрбиль, Ирак

## Решение интегральных уравнений Вольтерра-Фредгольма с помощью естественной кубической сплайн-функции

В статье с использованием функции натурального кубического сплайна найдено численное решение смешанных интегральных уравнений Вольтерра-Фредгольма второго рода. Предлагаемый метод основан на применении естественной кубической сплайн-функции неизвестной функции в произвольной точке и метода интегрирования для преобразования интегрального уравнения Вольтерра-Фредгольма в систему линейных уравнений относительно неизвестной функции. Приближенное решение легко получить, решив данную систему. Это достигается с помощью компьютерной программы, работающей на Python 3.9.

*Ключевые слова:* интегральное уравнение Вольтерра, интегральное уравнение Фредгольма, сплайн-функция.