

Asymptotic estimates of the solution for a singularly perturbed Cauchy problem

N.U. Bukanay^{1,*}, A.E. Mirzakulova¹, A.T. Assanova²

¹*al-Farabi Kazakh National University, Almaty, Kazakhstan;*

²*Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan
(E-mail: nbukanay@gmail.com, aziza.mirzakulova@bk.ru, anartasan@gmail.com)*

The article focuses on the initial problem for a third-order linear integro-differential equation with a small parameter at the higher derivatives, assuming that the roots of the additional characteristic equation have opposite signs. This paper presents a fundamental set of solutions and initial functions for a singularly perturbed homogeneous differential equation. The solution to the singularly perturbed initial integro-differential problem employs analytical formulas. A theorem concerning asymptotic estimates of the solution is established.

Keywords: singularly perturbed integro-differential equation, asymptotic estimates, Cauchy functions, fundamental solutions, small parameter.

2020 Mathematics Subject Classification: 34D15, 34E10, 34K26.

Introduction

Vasil'eva A.B. and Butuzov V.F. introduced the theory of singularly perturbed equations in their work [1]. Kassymov K.A. [2] investigated the most common cases of the Cauchy problem for singularly perturbed nonlinear systems of ordinary differential and integro-differential equations, as well as partial differential equations of hyperbolic type. Subsequently, singularly perturbed initial and boundary value problems with initial jumps were studied in [3,4]. Mirzakulova A.E. [5] extensively examines boundary value problems, particularly when the roots of the additional characteristic equation have opposing signs. While linear integro-differential equations are presented in [6], numerous papers have been dedicated to singularly perturbed integro-differential equations [7–12]. This article also provides an asymptotic solution for a singularly perturbed differential equation in a boundary value problem where the roots of the characteristic equation are opposite [13]. In addition, in recent years, significant work has been done on the numerical solution of integro-differential problems [14–17].

In this paper, we consider the initial problem for third-order linear integral-differential equations with a small parameter, where the roots of the corresponding characteristic equation have opposite signs. It is well known that there is no solution to a third-order linear differential equation with a small parameter (where the roots of the corresponding characteristic equation have opposite signs). However, we demonstrate that adding an integral term to the right-hand side yields an asymptotic formula for the solution. This article presents the findings of this research, which include an analytical formula and asymptotic estimates for solving a singularly perturbed integral-differential equation with a small parameter and initial conditions. Furthermore, a theorem on asymptotic estimation of these equations' solutions is proven.

*Corresponding author. E-mail: nbukanay@gmail.com

This research was funded by the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP23488301).

Received: 11 November 2024; Accepted: 26 March 2025.

© 2025 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1 Statement of the problem

Consider the singularly perturbed integro-differential equation with a small parameter

$$L_\varepsilon y \equiv \varepsilon^2 y''' + \varepsilon A_2(t)y'' + A_1(t)y' + A_0(t)y = F(t) + \int_0^1 \sum_{i=0}^2 H_i(t, x)y^{(i)}(x, \varepsilon) dx, \quad (1)$$

in the initial conditions

$$y^{(i)}(0, \varepsilon) = \alpha_i, \quad i = \overline{0, 2}. \quad (2)$$

ε is a small parameter, while α_i are known constants.

Assume the following conditions hold:

I. Functions $A_i(t)$, $i = \overline{0, 2}$ and $F(t)$ are continuously differentiable on a segment $0 \leq t \leq 1$, and functions $H_i(t, x)$, $i = \overline{0, 2}$ are continuously differentiable on a domain $D = \{0 \leq t \leq 1, 0 \leq x \leq 1\}$.

II. Roots of “additional characteristic equation” $\mu^2 + A_2(t)\mu + A_1(t) = 0$ satisfy the following inequalities $\mu_1(t) \neq \mu_2(t)$ and $\mu_1(t) < -\gamma_1 < 0$, $\mu_2(t) > \gamma_2 > 0$.

To construct the solution, we first identify auxiliary functions. This article [13] serves as a reference. From the third formula to the seventh, we obtained the required results.

We look for the solution to equation (1) with conditions (2) in the form

$$y(t, \varepsilon) = \sum_{i=1}^3 C_i y_i(t, \varepsilon) + \frac{1}{\varepsilon^2} \int_0^t K_0(t, s, \varepsilon) z(s, \varepsilon) ds - \frac{1}{\varepsilon^2} \int_t^1 K_1(t, s, \varepsilon) z(s, \varepsilon) ds; \quad (3)$$

here the fundamental solutions are $y_i(t, \varepsilon)$, $i = \overline{1, 3}$. $K_0(t, s, \varepsilon)$, $K_1(t, s, \varepsilon)$ are auxiliary functions expressed by article [13], C_i , $i = \overline{1, 3}$ are unknown constants.

We now denote the right side of equation (1) as follows

$$z(t, \varepsilon) = F(t) + \int_0^1 \sum_{i=0}^2 H_i(t, x)y^{(i)}(x, \varepsilon) dx.$$

Instead of $y(x, \varepsilon)$, we apply formula (3) to obtain the formula for $z(t, \varepsilon)$

$$\begin{aligned} z(t, \varepsilon) = f(t, \varepsilon) + \frac{1}{\varepsilon^2} \int_0^1 \left[\int_s^1 \left(\sum_{i=0}^2 H_i(t, x) K_0^{(i)}(x, s, \varepsilon) dx \right) z(s, \varepsilon) ds - \right. \\ \left. - \frac{1}{\varepsilon^2} \int_0^s \left(\sum_{i=0}^2 H_i(t, x) K_1^{(i)}(x, s, \varepsilon) dx \right) z(s, \varepsilon) ds, \right. \end{aligned}$$

here

$$\begin{aligned} f(t, \varepsilon) = F(t) + C_1 \int_0^1 \left(\sum_{i=0}^2 H_i(t, x) y_1(x, \varepsilon) \right) dx + C_2 \int_0^1 \left(\sum_{i=0}^2 H_i(t, x) y_2(x, \varepsilon) \right) dx + \\ + C_3 \int_0^1 \left(\sum_{i=0}^2 H_i(t, x) y_3(x, \varepsilon) \right) dx = F(t) + C_1 \phi_1(t, \varepsilon) + C_2 \phi_2(t, \varepsilon) + C_3 \phi_3(t, \varepsilon), \end{aligned}$$

$$\phi_i(t, \varepsilon) = \int_0^1 \left(H_0(t, x) y_i(x, \varepsilon) + H_1(t, x) y_i'(x, \varepsilon) + H_2(t, x) y_i''(x, \varepsilon) \right) dx, \quad i = \overline{1, 3}. \quad (4)$$

Now, we provide the following notation

$$H(t, s, \varepsilon) = \frac{1}{\varepsilon^2} \int_s^1 \left(\sum_{i=0}^2 H_i(t, x) K_0^{(i)}(x, s, \varepsilon) \right) dx - \frac{1}{\varepsilon^2} \int_0^s \left(\sum_{i=0}^2 H_i(t, x) K_1^{(i)}(x, s, \varepsilon) \right) dx.$$

III. The kernel $H(t, s, \varepsilon)$ does not have an eigenvalue with the value 1.

We get the Fredholm equation of the second type

$$z(t, \varepsilon) = f(t, \varepsilon) + \int_0^1 H(t, s, \varepsilon) z(s, \varepsilon) ds. \quad (5)$$

If condition III is satisfied, the solution of integral equation (5) is the only one and it is written

$$z(t, \varepsilon) = f(t, \varepsilon) + \int_0^1 R(t, s, \varepsilon) z(s, \varepsilon) ds, \quad (6)$$

where $R(t, s, \varepsilon)$ represents the resolvent of the kernel $H(t, s, \varepsilon)$.

Formula (6) is substituted for $z(s, \varepsilon)$ in formula (3), then formula (3) is written

$$\begin{aligned} y(t, \varepsilon) = & \sum_{i=1}^3 C_i y_i(t, \varepsilon) + \frac{1}{\varepsilon^2} \int_0^t K_0(t, s, \varepsilon) \left[F(s) + \sum_{i=1}^3 C_i \phi_i(t, \varepsilon) + \int_0^1 R(s, p, \varepsilon) \left(F(p) + \sum_{i=1}^3 C_i \phi_i(p, \varepsilon) \right) dp \right] ds - \\ & - \frac{1}{\varepsilon^2} \int_t^1 K_1(t, s, \varepsilon) \left[F(s) + \sum_{i=1}^3 C_i \phi_i(t, \varepsilon) + \int_0^1 R(s, p, \varepsilon) \left(F(p) + \sum_{i=1}^3 C_i \phi_i(p, \varepsilon) \right) dp \right] ds. \end{aligned}$$

Now, we provide the following notations:

$$Q_i(t, \varepsilon) = y_i(t, \varepsilon) + \frac{1}{\varepsilon^2} \int_0^t K_0(t, s, \varepsilon) \overline{\phi}_i(s, \varepsilon) ds - \frac{1}{\varepsilon^2} \int_t^1 K_1(t, s, \varepsilon) \overline{\phi}_i(s, \varepsilon) ds, \quad i = \overline{1, 3}, \quad (7)$$

$$P(t, \varepsilon) = \frac{1}{\varepsilon^2} \int_0^t K_0(t, s, \varepsilon) \overline{F}(s, \varepsilon) ds - \frac{1}{\varepsilon^2} \int_t^1 K_1(t, s, \varepsilon) \overline{F}(s, \varepsilon) ds, \quad (8)$$

then the formulas for $y(t, \varepsilon)$ are abbreviated

$$y^{(j)}(t, \varepsilon) = C_1 Q_1^{(j)}(t, \varepsilon) + C_2 Q_2^{(j)}(t, \varepsilon) + C_3 Q_3^{(j)}(t, \varepsilon) + P^{(j)}(t, \varepsilon), \quad j = \overline{0, 2}. \quad (9)$$

Substituting equation (9) into condition (2) yields the algebraic system for calculating C_i , where $i = \overline{1, 3}$:

$$\begin{cases} C_1 Q_1(0, \varepsilon) + C_2 Q_2(0, \varepsilon) + C_3 Q_3(0, \varepsilon) + P(0, \varepsilon) = \alpha_0, \\ C_1 Q_1'(0, \varepsilon) + C_2 Q_2'(0, \varepsilon) + C_3 Q_3'(0, \varepsilon) + P'(0, \varepsilon) = \alpha_1, \\ C_1 Q_1''(0, \varepsilon) + C_2 Q_2''(0, \varepsilon) + C_3 Q_3''(0, \varepsilon) + P''(0, \varepsilon) = \alpha_2. \end{cases} \quad (10)$$

As $\varepsilon \rightarrow 0$, the asymptotic behavior of C_i , $i = \overline{1, 3}$, is obtained when the main determinant of system (10) is $\delta(\varepsilon) \neq 0$.

$$\begin{aligned} C_1(\varepsilon) &= -\varepsilon \frac{\mu_2(0)(\alpha_0 y'_{30}(0) - \alpha_1)}{\mu_1(0)(\mu_2(0) - \mu_1(0))} + O(\varepsilon^2), \\ C_2(\varepsilon) &= -\varepsilon \frac{\alpha_0}{\phi_2(0)} \left(\overline{\phi_3}(0) + y'_{30}(0) \mu_1(0) \mu_2(0) - \right. \\ &\quad \left. - y'_{30}(0) \frac{\mu_2(0) \overline{\phi_1}(0)}{\mu_1(0)(\mu_2(0) - \mu_1(0))} \right) + \varepsilon \frac{\alpha_1}{\phi_2(0)} \left(\mu_1(0) \mu_2(0) - \right. \\ &\quad \left. - \frac{\mu_2(0) \overline{\phi_1}(0)}{\mu_1(0)(\mu_2(0) - \mu_1(0))} \right) - \varepsilon \frac{\overline{F}(0)}{\phi_2(0)} + O(\varepsilon^2), \\ C_3(\varepsilon) &= \alpha_0 + O(\varepsilon). \end{aligned} \quad (11)$$

By formula (4) $\phi_i(t, \varepsilon)$, $i = \overline{1, 3}$ are found

$$\begin{aligned} \overline{\phi_1}(s, \varepsilon) &= \frac{1}{\varepsilon} \left(-\overline{H}_2(s, 0) \mu_1(0) y_{10}(0) + O(\varepsilon) \right) = \frac{1}{\varepsilon} \left(\overline{\phi_1}(s) + O(\varepsilon) \right), \\ \overline{\phi_2}(s, \varepsilon) &= \frac{1}{\varepsilon} \left(\overline{H}_2(s, 1) \mu_2(1) y_{20}(1) + O(\varepsilon) \right) = \frac{1}{\varepsilon} \left(\overline{\phi_2}(s) + O(\varepsilon) \right), \\ \overline{\phi_3}(s, \varepsilon) &= \int_0^1 \sum_{j=0}^2 \overline{H}_j(s, x) y_{30}^{(j)}(x) dx + O(\varepsilon) = \overline{\phi_3}(s) + O(\varepsilon). \end{aligned}$$

Given formulas (7) and (8), we obtain the asymptotic behaviors of $Q_i^{(j)}(t, \varepsilon)$ and $P^{(j)}(t, \varepsilon)$, $j = \overline{0, 2}$, $i = \overline{1, 3}$:

$$\begin{aligned} Q_1^{(j)}(t, \varepsilon) &= \frac{1}{\varepsilon} \int_0^t \frac{y_{30}^{(j)}(s) \overline{\phi_1}(s)}{\mu_1(s) \mu_2(s) y_{30}(s)} ds + \frac{1}{\varepsilon^j} e^{\frac{1}{\varepsilon} \int_0^t \mu_1(x) dx} \left(y_{10}(t) \mu_1^j(t) - \frac{\overline{\phi_1}(0) y_{10}(t) \mu_1^j(t)}{y_{10}(0) \mu_1^2(0) (\mu_2(0) - \mu_1(0))} \right) + \\ &+ \frac{1}{\varepsilon^j} e^{-\frac{1}{\varepsilon} \int_t^1 \mu_2(x) dx} \frac{\overline{\phi_1}(1) y_{20}(t) \mu_2^j(t)}{y_{20}(1) \mu_2^2(1) (\mu_2(1) - \mu_1(1))} + \frac{1}{\varepsilon^j} \frac{\overline{\phi_1}(t)}{(\mu_2(t) - \mu_1(t))} \left(\frac{\mu_1^j(t)}{\mu_1^2(t)} - \frac{\mu_2^j(t)}{\mu_2^2(t)} \right), \quad j = \overline{0, 2}, \quad (12.a) \end{aligned}$$

$$\begin{aligned} Q_2^{(j)}(t, \varepsilon) &= \frac{1}{\varepsilon} \int_0^t \frac{y_{30}^{(j)}(s) \overline{\phi_2}(s)}{\mu_1(s) \mu_2(s) y_{30}(s)} ds - \frac{1}{\varepsilon^j} e^{\frac{1}{\varepsilon} \int_0^t \mu_1(x) dx} \frac{\overline{\phi_2}(0) y_{10}(t) \mu_1^j(t)}{y_{10}(0) \mu_1^2(0) (\mu_2(0) - \mu_1(0))} + \\ &+ \frac{1}{\varepsilon^j} e^{-\frac{1}{\varepsilon} \int_t^1 \mu_2(x) dx} \left(y_{20}(t) \mu_2^{(j)}(t) + \frac{\overline{\phi_2}(1) y_{20}(t) \mu_2^j(t)}{y_{20}(1) \mu_2^2(1) (\mu_2(1) - \mu_1(1))} \right) + \\ &+ \frac{1}{\varepsilon^j} \frac{\overline{\phi_2}(t)}{(\mu_2(t) - \mu_1(t))} \left(\frac{\mu_1^j(t)}{\mu_1^2(t)} - \frac{\mu_2^j(t)}{\mu_2^2(t)} \right), \quad j = \overline{0, 2}, \quad (12.b) \end{aligned}$$

$$Q_3^{(j)}(t, \varepsilon) = y_{30}^{(j)}(t) + \int_0^t \frac{y_{30}^{(j)}(s) \overline{\phi_3}(s)}{\mu_1(s) \mu_2(s) y_{30}(s)} ds - \frac{1}{\varepsilon^{j-1}} e^{\frac{1}{\varepsilon} \int_0^t \mu_1(x) dx} \frac{\overline{\phi_3}(0) y_{10}(t) \mu_1^j(t)}{y_{10}(0) \mu_1^2(0) (\mu_2(0) - \mu_1(0))} +$$

$$+ \frac{1}{\varepsilon^{j-1}} e^{-\frac{1}{\varepsilon} \int_t^1 \mu_2(x) dx} \frac{\bar{\phi}_3(1) y_{20}(t) \mu_2^j(t)}{y_{20}(1) \mu_2^2(1) (\mu_2(1) - \mu_1(1))} + \frac{1}{\varepsilon^{j-1}} \frac{\bar{\phi}_3(t)}{(\mu_2(t) - \mu_1(t))} \left(\frac{\mu_1^j(t)}{\mu_1^2(t)} - \frac{\mu_2^j(t)}{\mu_2^2(t)} \right), \quad j = \overline{0, 2}, \quad (12.c)$$

$$P^{(j)}(t, \varepsilon) = \int_0^t \frac{y_{30}^{(j)}(s) \bar{F}(s)}{\mu_1(s) \mu_2(s) y_{30}(s)} ds - \frac{1}{\varepsilon^{j-1}} e^{\frac{1}{\varepsilon} \int_0^t \mu_1(x) dx} \frac{\bar{F}(0) y_{10}(t) \mu_1^j(t)}{y_{10}(0) \mu_1^2(0) (\mu_2(0) - \mu_1(0))} +$$

$$+ \frac{1}{\varepsilon^{j-1}} e^{-\frac{1}{\varepsilon} \int_t^1 \mu_2(x) dx} \frac{\bar{F}(s) y_{20}(t) \mu_2^j(t)}{y_{20}(1) \mu_2^2(1) (\mu_2(1) - \mu_1(1))} + \frac{1}{\varepsilon^{j-1}} \frac{\bar{F}(t)}{(\mu_2(t) - \mu_1(t))} \left(\frac{\mu_1^j(t)}{\mu_1^2(t)} - \frac{\mu_2^j(t)}{\mu_2^2(t)} \right), \quad j = \overline{0, 2}. \quad (12.d)$$

Theorem 1. If conditions I-III are valid, then the solution for integro-differential equation (1) and (2) holds the following asymptotic estimates as $\varepsilon \rightarrow 0$:

$$|y^{(j)}(t, \varepsilon)| \leq C \left(|\alpha_0| + \varepsilon |\alpha_1| + \varepsilon^2 |\alpha_2| + \max_{0 \leq t \leq 1} |F(t)| \right) +$$

$$+ \frac{C}{\varepsilon^{j-1}} e^{-\frac{t}{\varepsilon} \gamma_1} \left(|\alpha_0| + |\alpha_1| + \varepsilon^2 |\alpha_2| + \max_{0 \leq t \leq 1} |F(t)| \right) + \quad (13)$$

$$+ \frac{C}{\varepsilon^{j-1}} e^{-\frac{1-t}{\varepsilon} \gamma_2} \left(|\alpha_0| + |\alpha_1| + \varepsilon^2 |\alpha_2| + \max_{0 \leq t \leq 1} |F(t)| \right), \quad j = 0, 1, 2,$$

where $C > 0$ is a constant independent of ε .

Proof. Asymptotic estimates of C_i , $i = \overline{1, 3}$ and $Q_i^{(j)}(t, \varepsilon)$, $P^{(j)}(t, \varepsilon)$ are obtained by applying formulas (11)-(12.a-12.d):

$$|C_i| \leq C \varepsilon \left(|\alpha_0| + |\alpha_1| \right), \quad i = 1, 2,$$

$$|C_3| \leq C \left(|\alpha_0| + \varepsilon^2 |\alpha_2| \right),$$

$$|Q_1^{(j)}(t, \varepsilon)| \leq C \left(1 + \frac{1}{\varepsilon^j} e^{-\frac{t}{\varepsilon} \gamma_1} + \frac{1}{\varepsilon^j} e^{-\frac{1-t}{\varepsilon} \gamma_2} \right), \quad j = \overline{0, 2},$$

$$|Q_2^{(j)}(t, \varepsilon)| \leq C \left(1 + \frac{1}{\varepsilon^j} e^{-\frac{t}{\varepsilon} \gamma_1} + \frac{1}{\varepsilon^j} e^{-\frac{1-t}{\varepsilon} \gamma_2} \right), \quad j = \overline{0, 2}, \quad (14)$$

$$|Q_3^{(j)}(t, \varepsilon)| \leq C \left(1 + \frac{1}{\varepsilon^{j-1}} e^{-\frac{t}{\varepsilon} \gamma_1} + \frac{1}{\varepsilon^{j-1}} e^{-\frac{1-t}{\varepsilon} \gamma_2} \right), \quad j = \overline{0, 2},$$

$$|P^{(j)}(t, \varepsilon)| \leq C \max_{0 \leq t \leq 1} |F(t)| \left(1 + \frac{1}{\varepsilon^{j-1}} e^{-\frac{t}{\varepsilon} \gamma_1} + \frac{1}{\varepsilon^{j-1}} e^{-\frac{1-t}{\varepsilon} \gamma_2} \right), \quad j = \overline{0, 2}.$$

We derive asymptotic estimations (13) from asymptotic behaviors (14). Theorem 1 has been proved.

Conclusion

This article examines the initial problem for a third-order linear integro-differential equation with a small parameter at the higher derivatives, assuming that the roots of the additional characteristic equation have opposite signs. This paper presents the construction of a fundamental system of solutions and a Cauchy function for a singularly perturbed homogeneous differential equation. The functions $Q_i(t, \varepsilon)$, $P(t, \varepsilon)$, $i = \overline{1, 3}$, and constants C_i , $i = \overline{1, 3}$ exhibit asymptotic behaviors and estimates. Furthermore, the article provides an analytical formula for solving this singularly perturbed initial problem. A theorem on asymptotic estimates of the solution is proven.

Acknowledgments

The present work was partially supported by the Grant “Asymptotic estimates of solutions of singularly perturbed differential, integro-differential and impulse equations” (2024–2026) of the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP23488301).

Author Contributions

N.U. Bukanay used the fundamental solutions of the homogeneous equation to calculate the Cauchy function and derive the asymptotic formula. A.E. Mirzakulova checked all the calculations and wrote the asymptotic behavior of the solutions. A.T. Assanova conducted a comprehensive verification of the calculations. All authors contributed equally to this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Vasil'eva, A.B., & Butuzov, V.F. (1973). *Asimptoticheskie razlozheniia reshenii singuliarno vozmushchennykh uravnenii* [Asymptotic expansions of solutions of singularly perturbed equations]. Moscow: Nauka [in Russian].
- 2 Kassymov, K.A. (1987). *Singuliarno vozmushchennye kraevye zadachi s nachalnymi skachkami* [Singularly perturbed boundary value problems with initial jumps]. Almaty: Izdatelstvo “Sanat” [in Russian].
- 3 Kassymov, K.A., & Dauylbayev, M.K. (1999). Estimates of solutions of the Cauchy problem with an arbitrary-order initial jump for linear singularly perturbed integro-differential equations. *Differential Equations*, 35(6), 822–830.
- 4 Nurgabul, D. (2014). Asymptotic estimates for the Solution of a Restoration Problem with Initial Jump. *Journal of Applied Mathematic*, 2014, 956402.
- 5 Mirzakulova, A.E. (2023). *Kishi parametrleri bar tendeler ushin bastapqy zhane shettik esep-terdi sheshu adisteri* [Methods for solving initial and boundary problems for equations with small parameters]. (Vol. 176). Almaty: Qazaq universiteti [in Kazakh].
- 6 Dauylbayev, M.K. (2009). *Lineinye integro-differentsialnye uravneniia s malym parametrom* [Linear integro-differential equations with a small parameter]. Almaty: Qazaq universiteti [in Russian].

- 7 Bobodzhanova, M.A., Kalimbetov, B.T., & Safonov, V.F. (2024). Singularly perturbed integro-differential equations with degenerate Hammerstein's kernel. *Bulletin of the Karaganda University. Mathematics Series*, 4(116), 57–68. <https://doi.org/10.31489/2024M4/57-68>
- 8 Kellogg, R.B., & Stynes, M. (2006). A singularly perturbed convection-diffusion problem in a half-plane. *Appl. Anal.*, 85(12), 1471–1485. <http://dx.doi.org/10.1080/00036810601066574>
- 9 Gracia, J.L., & O'Riordan, E. (2012). A singularly perturbed parabolic problem with a layer in the initial condition. *Appl. Math. Comput.*, 219(2), 498–510. <http://dx.doi.org/10.1016/j.amc.2012.06.028>
- 10 Assanova, A.T. (2021). Hyperbolic equation piecewise-constant argument of generalized type and solving boundary value problems for it. *Lobachevskii Journal of Mathematics*, 42(15), 3584–3593. <http://dx.doi.org/10.1134/S1995080222030040>
- 11 Oljira, A.F., & Woldaregay, M.M. (2024). A fitted operator numerical method for singularly perturbed Fredholm integro-differential equation with integral initial condition. *BMC Research Notes*, 17(1), 23. <http://dx.doi.org/10.1186/s13104-023-06649-9>
- 12 Tiruneh, A.A., Kumie, H.G., & Derese, G.A. (2024). Singularly perturbed time-fractional convection–diffusion equations via exponential fitted operator scheme. *Partial Differential Equations in Applied Mathematics*, 11(1), 100873. <http://dx.doi.org/10.1016/j.padiff.2024.100873>
- 13 Bukanay, N.U., Mirzakulova, A.E., Dauylbayev, M.K., & Konisbayeva, K.T. (2020). A boundary jumps phenomenon in the integral boundary value problem for singularly perturbed differential equations. *Bulletin of the Karaganda University. Mathematics Series*, 2(98), 46–58. <https://doi.org/10.31489/2020M2/46-58>
- 14 Oljira, A.F., & Woldaregay, M.M. (2024). A fitted operator numerical method for singularly perturbed Fredholm integro-differential equation with integral initial condition. *BMC Research Notes*, 17, 23. <https://doi.org/10.1186/s13104-023-06649-9>
- 15 Durmaz, M.E., Cakir, M., Amirali, I., & Amiraliyev, G.M. (2022). Numerical solution of singularly perturbed Fredholm integro-differential equations by homogeneous second order difference method. *Journal of Computational and Applied Mathematics*, 412(1), 114327. <https://doi.org/10.1016/j.cam.2022.114327>
- 16 Rathore, A.S., & Shanthi, V. (2024). A numerical solution of singularly perturbed Fredholm integro-differential equation with discontinuous source term. *Journal of Computational and Applied Mathematics*, 446, 115858. <https://doi.org/10.1016/j.cam.2024.115858>
- 17 Panda, A., Mohapatra, J., Amirali, I., Durmaz, M.E., & Amiraliyev, G.M. (2024). A numerical technique for solving nonlinear singularly perturbed Fredholm integro-differential equations. *Mathematics and Computers in Simulation*, 220, 618–629. <https://doi.org/10.1016/j.matcom.2024.02.011>

*Author Information**

Nuraiym Ulankyzy Bukanay — Second-year PhD student, Teacher, Department of Mathematics, al-Farabi Kazakh National University, 71 al-Farabi Avenue, Almaty, 050038, Kazakhstan; e-mail: nbukanay@gmail.com; <https://orcid.org/0009-0009-2206-2302>

Aziza Erkomekovna Mirzakulova — Acting Associate Professor, PhD, Department of Mathematics, al-Farabi Kazakh National University, 71 al-Farabi Avenue, Almaty, 050038, Kazakhstan; e-mail: aziza.mirzakulova@bk.ru; <https://orcid.org/0000-0001-6445-6371>

*The author's name is presented in the order: First, Middle, and Last Names.

Anar Turmaganbetkyzy Assanova — Dr. of Sciences in Phys. and Math., Professor, Institute of Mathematics and Mathematical Modeling, Senior Researcher, Almaty, Kazakhstan; e-mail: *anartasan@gmail.com*; <http://orcid.org/0000-0001-8697-8920>